PROCESS OF OBTAINING YOUNG AGAVE FRUCTANS AND USE AS A FUNCTIONAL INGREDIENT		
Offering Organization:	Centro de Investigación y Asistencia en Tecnología y Diseño del Estado	
	de Jalisco, A.C.	
Type of Organization:	Public Research Center	
Dovelopment Stage:	Laboratory	
Development Stage: Desired Relationship:	Laboratory	
Desirea keiationship.	 Technological research and development financing (technological partner) 	
	 Specialized application tests Creation of a new company (Joint Venture) for the 	
	commercialization of the products outlined herein	
	 Licensing of patents 	
Sector:	Food	
Area of knowledge:	Food Technology	
Key words:	Fructans of young agave; Nutraceuticals	
DETAILED DESCRIPTION:		
Problem to be solved:		
The processing and use of agave fructans as a functional ingredient with nutraceutical		
properties to reduce serum triglycerides in overweight or obese patients, or in normal-		
	hat suffer from dyslipidemia.	
Solution:	,	
A product that	contains a mixture of agave fructans with different degrees of	
-	which can be used as a nutraceutical ingredient in the formulation of	
functional foods	or for use as a food supplement. This product reduces triglycerides that	
are present in me	etabolic disorders and overweight or obese persons.	
are present in me		
New and Innovative Aspe	etabolic disorders and overweight or obese persons.	
New and Innovative Aspe The agave used c	etabolic disorders and overweight or obese persons. <i>Acts:</i> letermines the differentiation of its carbohydrates. This is due to the fact	
New and Innovative Aspe The agave used c	etabolic disorders and overweight or obese persons.	
New and Innovative Aspe The agave used o that the polymer	etabolic disorders and overweight or obese persons. <i>acts:</i> letermines the differentiation of its carbohydrates. This is due to the fact chains containing agave fructans increase in size over time.	
New and Innovative Aspe The agave used c	etabolic disorders and overweight or obese persons. <i>acts:</i> letermines the differentiation of its carbohydrates. This is due to the fact chains containing agave fructans increase in size over time.	
New and Innovative Aspe The agave used o that the polymer	etabolic disorders and overweight or obese persons. <i>Acts:</i> letermines the differentiation of its carbohydrates. This is due to the fact chains containing agave fructans increase in size over time. STICS:	
New and Innovative Aspe The agave used o that the polymer TECHNICAL CHARACTERI This product is	etabolic disorders and overweight or obese persons. Acts: letermines the differentiation of its carbohydrates. This is due to the fact chains containing agave fructans increase in size over time. STICS: formed from a mixture of agave fructans with different degrees of	
New and Innovative Aspe The agave used o that the polymer TECHNICAL CHARACTERI This product is polymerization. T	etabolic disorders and overweight or obese persons. Acts: letermines the differentiation of its carbohydrates. This is due to the fact chains containing agave fructans increase in size over time. STICS: formed from a mixture of agave fructans with different degrees of These are specifically found in agave plants with a maturity of 3 years. It	
New and Innovative Aspe The agave used of that the polymer TECHNICAL CHARACTERI This product is polymerization. T is composed of	etabolic disorders and overweight or obese persons. <i>Acts:</i> letermines the differentiation of its carbohydrates. This is due to the fact chains containing agave fructans increase in size over time. STICS: formed from a mixture of agave fructans with different degrees of These are specifically found in agave plants with a maturity of 3 years. It long/short chain fructans in ratios ranging from 30/70 to 50/50. This	
New and Innovative Aspe The agave used of that the polymer TECHNICAL CHARACTERI This product is polymerization. T is composed of product is derive	etabolic disorders and overweight or obese persons. Acts: letermines the differentiation of its carbohydrates. This is due to the fact chains containing agave fructans increase in size over time. STICS: formed from a mixture of agave fructans with different degrees of These are specifically found in agave plants with a maturity of 3 years. It long/short chain fructans in ratios ranging from 30/70 to 50/50. This d from young agave plants. The agave is washed and the fiber is torn off	
New and Innovative Aspe The agave used of that the polymer TECHNICAL CHARACTERI This product is polymerization. T is composed of product is derive the plant and gro	etabolic disorders and overweight or obese persons. Acts: letermines the differentiation of its carbohydrates. This is due to the fact chains containing agave fructans increase in size over time. STICS: formed from a mixture of agave fructans with different degrees of These are specifically found in agave plants with a maturity of 3 years. It long/short chain fructans in ratios ranging from 30/70 to 50/50. This d from young agave plants. The agave is washed and the fiber is torn off pund, and then combined with hot water to obtain an aqueous extract	
New and Innovative Aspe The agave used of that the polymer TECHNICAL CHARACTERI This product is polymerization. T is composed of product is derive the plant and gro containing agav	etabolic disorders and overweight or obese persons. Acts: letermines the differentiation of its carbohydrates. This is due to the fact chains containing agave fructans increase in size over time. STICS: formed from a mixture of agave fructans with different degrees of These are specifically found in agave plants with a maturity of 3 years. It long/short chain fructans in ratios ranging from 30/70 to 50/50. This d from young agave plants. The agave is washed and the fiber is torn off bund, and then combined with hot water to obtain an aqueous extract e soluble solids. The extract is then filtered using coagulation,	
New and Innovative Aspe The agave used of that the polymer TECHNICAL CHARACTERI This product is polymerization. T is composed of product is derive the plant and gro containing agav sedimentation a	etabolic disorders and overweight or obese persons. Acts: letermines the differentiation of its carbohydrates. This is due to the fact chains containing agave fructans increase in size over time. STICS: formed from a mixture of agave fructans with different degrees of These are specifically found in agave plants with a maturity of 3 years. It long/short chain fructans in ratios ranging from 30/70 to 50/50. This d from young agave plants. The agave is washed and the fiber is torn off pund, and then combined with hot water to obtain an aqueous extract	
New and Innovative Aspe The agave used of that the polymer TECHNICAL CHARACTERI This product is polymerization. T is composed of product is derive the plant and gro containing agav sedimentation and can be concentra	etabolic disorders and overweight or obese persons. Acts: letermines the differentiation of its carbohydrates. This is due to the fact chains containing agave fructans increase in size over time. STICS: formed from a mixture of agave fructans with different degrees of These are specifically found in agave plants with a maturity of 3 years. It long/short chain fructans in ratios ranging from 30/70 to 50/50. This d from young agave plants. The agave is washed and the fiber is torn off pund, and then combined with hot water to obtain an aqueous extract e soluble solids. The extract is then filtered using coagulation, nd filtration techniques to remove insoluble solids present. The extract ted or dehydrated to obtain a powder.	
New and Innovative Aspe The agave used of that the polymer TECHNICAL CHARACTERI This product is polymerization. This composed of product is derive the plant and gro containing agav sedimentation and can be concentration	etabolic disorders and overweight or obese persons. Acts: letermines the differentiation of its carbohydrates. This is due to the fact chains containing agave fructans increase in size over time. STICS: formed from a mixture of agave fructans with different degrees of These are specifically found in agave plants with a maturity of 3 years. It long/short chain fructans in ratios ranging from 30/70 to 50/50. This d from young agave plants. The agave is washed and the fiber is torn off pund, and then combined with hot water to obtain an aqueous extract e soluble solids. The extract is then filtered using coagulation, nd filtration techniques to remove insoluble solids present. The extract ted or dehydrated to obtain a powder. I from its utilization:	
New and Innovative Aspe The agave used of that the polymer TECHNICAL CHARACTERI This product is polymerization. T is composed of product is derive the plant and gro containing agav sedimentation an can be concentra Main advantages derived – This product	etabolic disorders and overweight or obese persons. Acts: letermines the differentiation of its carbohydrates. This is due to the fact chains containing agave fructans increase in size over time. STICS: formed from a mixture of agave fructans with different degrees of These are specifically found in agave plants with a maturity of 3 years. It long/short chain fructans in ratios ranging from 30/70 to 50/50. This d from young agave plants. The agave is washed and the fiber is torn off bund, and then combined with hot water to obtain an aqueous extract e soluble solids. The extract is then filtered using coagulation, nd filtration techniques to remove insoluble solids present. The extract ted or dehydrated to obtain a powder. <i>I from its utilization:</i> is a mixture of branched agave fructans, it contains a higher proportion	
New and Innovative Aspective The agave used of that the polymer TECHNICAL CHARACTERI This product is polymerization. This composed of product is derived the plant and grad containing agav sedimentation and can be concentrated Main advantages derived of short-chai	etabolic disorders and overweight or obese persons. Acts: letermines the differentiation of its carbohydrates. This is due to the fact chains containing agave fructans increase in size over time. STICS: formed from a mixture of agave fructans with different degrees of These are specifically found in agave plants with a maturity of 3 years. It long/short chain fructans in ratios ranging from 30/70 to 50/50. This d from young agave plants. The agave is washed and the fiber is torn off bund, and then combined with hot water to obtain an aqueous extract e soluble solids. The extract is then filtered using coagulation, nd filtration techniques to remove insoluble solids present. The extract ted or dehydrated to obtain a powder. <i>I from its utilization:</i> is a mixture of branched agave fructans, it contains a higher proportion in fructans with degrees of polymerizations of 3 to 10 fructose units	
New and Innovative Aspe The agave used of that the polymer TECHNICAL CHARACTERI This product is polymerization. T is composed of product is derive the plant and gro containing agav sedimentation an can be concentra Main advantages derived of short-chai – This product	etabolic disorders and overweight or obese persons. Acts: Letermines the differentiation of its carbohydrates. This is due to the fact chains containing agave fructans increase in size over time. STICS: formed from a mixture of agave fructans with different degrees of These are specifically found in agave plants with a maturity of 3 years. It long/short chain fructans in ratios ranging from 30/70 to 50/50. This d from young agave plants. The agave is washed and the fiber is torn off bund, and then combined with hot water to obtain an aqueous extract e soluble solids. The extract is then filtered using coagulation, nd filtration techniques to remove insoluble solids present. The extract ted or dehydrated to obtain a powder. I from its utilization: is a mixture of branched agave fructans, it contains a higher proportion in fructans with degrees of polymerizations of 3 to 10 fructose units is decreases the concentration of serum triglycerides, which can be	
New and Innovative Aspe The agave used of that the polymer TECHNICAL CHARACTERI This product is polymerization. T is composed of product is derive the plant and gro containing agav sedimentation an can be concentra Main advantages derived – This product of short-chai – This product	etabolic disorders and overweight or obese persons. Acts: letermines the differentiation of its carbohydrates. This is due to the fact chains containing agave fructans increase in size over time. STICS: formed from a mixture of agave fructans with different degrees of These are specifically found in agave plants with a maturity of 3 years. It long/short chain fructans in ratios ranging from 30/70 to 50/50. This d from young agave plants. The agave is washed and the fiber is torn off bund, and then combined with hot water to obtain an aqueous extract e soluble solids. The extract is then filtered using coagulation, nd filtration techniques to remove insoluble solids present. The extract ted or dehydrated to obtain a powder. <i>I from its utilization:</i> is a mixture of branched agave fructans, it contains a higher proportion in fructans with degrees of polymerizations of 3 to 10 fructose units	

Applications: – Food supplements, pharmaceutical components		
INTELLECTUAL PROPERTY		
 Patent application submitted in 2013 Application number : MX/a/2013/004901 		
ABOUT THE OFFERING ORGANIZATION		
Presentation:	El Centro de Investigación y Asistencia en Tecnología y Diseño del Estado de Jalisco, A.C. (CIATEJ) is a public research center that belongs to the national technology development and innovation network, the National Council for Science and Technology (CONACyT). CIATEJ is focused on the agricultural, food, health, and environmental sectors with an emphasis on the application of innovative biotechnology.	
Contact Information:	Evaristo Urzúa Esteva - <u>eurzua@ciatej.net.mx</u>	